

Neutrino textures and charged lepton flavour violation in light of θ_{13} , MEG and LHC data

M. Cannoni¹, J. Ellis^{2,3}, M.E. Gómez¹ and S. Lola^{*}

¹ Department of Applied Physics, University of Huelva, 21071 Huelva, Spain

² Theoretical Particle Physics and Cosmology Group, Department of Physics,
King's College London, Strand, London WC2R 2LS, UK

³ Theory Division, Physics Department, CERN, CH-1211 Geneva 23, Switzerland

ABSTRACT

In light of recent results from the LHC, MEG and neutrino experiments, we revisit the issue of charged lepton flavour violation (LFV), which arises naturally in supersymmetric theories with massive neutrinos via flavour-violating soft supersymmetry-breaking masses for sleptons induced by radiative corrections. We link our results to the expectations for neutrino mixing angles, with particular focus on θ_{13} in various Abelian flavour texture models, showing how LFV observables may distinguish between different possibilities. We focus on the radiative decays $\mu \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$, and on detection prospects at the LHC and a linear collider (LC). We use supersymmetric parameters favoured by $g_\mu - 2$ and cosmological considerations that are consistent with LHC searches for supersymmetry and the Higgs mass. We find a class of scenarios where the LHC may be sensitive to LFV sparticle decays and LFV processes could be detectable at a LC with centre-of-mass energy above 1 TeV, whereas LFV lepton decays may be suppressed by cancellations in the decay amplitudes.

^{*}On leave from the Department of Physics, University of Patras, 26500 Patras, Greece.

1 Introduction

In recent years, the existence of neutrino masses and oscillations with near-maximal $\nu_\mu - \nu_\tau$ and large $\nu_e \rightarrow \nu_\mu$ mixing has been established by extensive input from atmospheric [1], solar [2] and long-baseline reactor [3] and accelerator [4, 5] neutrino experiments. Initial input on the possible range of the θ_{13} [6, 7] was provided by the T2K [8] and MINOS collaborations [9], and definitive evidence for a non-zero value of θ_{13} has been provided by the reactor experiments Daya Bay [10] and RENO [11], and very recently also by Double Chooz [12].

A natural expectation in theories with massive neutrinos is charged-lepton-flavour violation (LFV), which is enhanced in supersymmetric theories via the renormalization of soft supersymmetry-breaking parameters. The link between neutrino oscillations and violations of the individual lepton numbers $L_{e,\mu,\tau}$ raises the prospect of observing processes such as $\mu \rightarrow e\gamma$, $\mu \rightarrow 3e$, $\tau \rightarrow \mu\gamma$ and $\mu \rightarrow e$ conversion on heavy nuclei [13].

The present experimental upper limits on the most interesting of these processes, summarised below, already constrain significantly the parameter spaces of theoretical models.

$$BR(\mu \rightarrow e\gamma) < 2.4 \times 10^{-12} \quad [14], \quad (1)$$

$$BR(\tau \rightarrow \mu\gamma) < 4.4 \times 10^{-8} \quad [15], \quad (2)$$

$$BR(\tau \rightarrow e\gamma) < 3.3 \times 10^{-8} \quad [15]. \quad (3)$$

The strongest constraint on radiative decays is the MEG upper limit on $BR(\mu \rightarrow e\gamma)$ [14].

Within the supersymmetric framework, one should also keep in mind other possibilities for observing LFV processes, such as slepton pair production at a Linear Collider (LC) [16, 17, 18, 19, 20, 21, 22] and signals at the LHC [23, 24, 25, 26, 27, 28, 29], particularly in $\chi_2 \rightarrow \chi + e^\pm \mu^\mp$, $\chi_2 \rightarrow \chi + \mu^\pm \tau^\mp$ decays (here χ is the lightest neutralino, assumed to be the lightest supersymmetric particle (LSP), and χ_2 is the second-lightest neutralino). These decays could provide search prospects that are complementary to direct searches for the flavour-violating decays of charged leptons, particularly for heavy superparticle spectra.

In this paper we re-evaluate the prospects for observable charged LFV, based on updated knowledge of neutrino mass and mixing parameters, including the recent measurement of θ_{13} . We work within the framework of the most natural mechanism for obtaining light neutrino masses, namely the see-saw mechanism [30], in which an effective Majorana mass matrix for light neutrinos, $m_{eff}^D = m_\nu^D \cdot (M_N)^{-1} \cdot m_\nu^{D^T}$, arises from Dirac neutrino masses m_ν^D of the same order as the charged-lepton and quark masses, and heavy Majorana masses M_N .

In supersymmetric theories, the neutrino Dirac couplings Y_ν renormalise the soft supersymmetry-breaking sneutrino and slepton masses, generating LFV naturally [31]. Even if the soft scalar masses were universal at the unification scale, quantum corrections between the GUT scale and low energies would modify this structure via renormalization-group running, which generates off-diagonal contributions. This effect is particularly interesting in see-saw models, where in general the Dirac neutrino Yukawa couplings cannot be diagonalized simultaneously with the charged-lepton and slepton mass matrices [31]. Given the large mixing of the corresponding neu-

trino species, charged-lepton-flavour violation may occur at enhanced rates in supersymmetric extensions of the standard model, giving rise to observable LFV signals [13, 32, 33, 34, 35, 36].

We analyse this possibility within the constrained minimal supersymmetric standard model (CMSSM) with universal scalar, gaugino masses and trilinear terms at the GUT scale, m_0 , $M_{1/2}$ and A_0 , respectively, using textures for mass matrices that are inspired by GUT models with Abelian flavour symmetries [37, 38]. These textures reproduce naturally the observed fermion mass hierarchies and mixing angles. Despite their phenomenological appeal, however, there are ambiguities and limitations due to the fact that the entries in the mass matrices are determined only up to $\mathcal{O}(1)$ numerical factors. In a recent paper analysing textures derived by an SU(5) Froggatt-Nielsen mechanism, where all coefficients were treated as random variables in a Monte-Carlo scan of the multi-dimensional parameter space, it was shown that values of θ_{13} matching the recent data are statistically favoured [39]. Additional work in the same context may also be found in Ref. [40].

In this work, we extend these studies, showing that the neutrino parameters favoured by the data offer good detection prospects for detection of charged LFV in several channels. We study the complementarity between different radiative decays, deriving specific information on the allowed correlations between the decay rates. In all cases, we focus on sparticle spectra and values of $\tan\beta$ that are compatible with searches at the LHC, and also compatible with cosmological relic density considerations [41]. As an output of our analysis, we discuss detection prospects at the LHC and a Linear Collider (LC), taking into account the constraints that the recent LHC supersymmetry searches and the measurement of the Higgs mass impose on the observability of LFV processes. As we discuss later in more detail, one of the scenarios favoured by these constraints has relatively large right-handed sparticle mixing, and hence may lead to observable LFV in cascade decays to sleptons at the LHC [26], as well as at a LC with a centre-of-mass energy above 1 TeV.

2 Neutrino Mass Textures inspired by SU(5)

Over the recent years, a plethora of textures have been proposed to explain the data on neutrino masses and mixing. The new data on θ_{13} provide additional constraints, excluding certain possibilities and constraining others. In our work, rather than reviewing the vast literature on the subject, we choose a representative model that fits the fermion data and is well-motivated on theoretical grounds. Nevertheless, we try to keep the results as generic as possible, placing emphasis on the links between physical observables. We also keep in mind that several *a priori* different theoretical models may converge to similar phenomenology, since they are matched to the same data.

The example we choose is provided by a SU(5) GUT combined with family symmetries [37, 38]. The mass matrices are constructed by looking at the field content of the SU(5) representations, namely: three families of $(Q, u^c, e^c)_i \in \mathbf{10}$ in representations, three families of $(L, d^c)_i \in \bar{\mathbf{5}}$ representations, and heavy right-handed neutrinos in singlet representations. This model therefore

has the following properties: (i) the up-quark mass matrix is symmetric, and (ii) the charged-lepton mass matrix is the transpose of the down-quark mass matrix, which relates the mixing of the left-handed leptons to that of the right-handed down-type quarks. Since the Cabibbo-Kobayashi-Maskawa (CKM) mixing in the quark sector is due to a mismatch between the mixing of the left-handed up- and down-type quarks, it is independent from the mixing in the lepton sector. In this way, it is easy to reconcile large atmospheric neutrino mixing with the observed small V_{CKM} mixing angles.

Within this framework, and following [37, 38, 42, 43], for example, the Yukawa matrices have the form

$$Y_u \propto \begin{pmatrix} \varepsilon^6 & \varepsilon^5 & \varepsilon^3 \\ \varepsilon^5 & \varepsilon^4 & \varepsilon^2 \\ \varepsilon^3 & \varepsilon^2 & 1 \end{pmatrix}, \quad Y_\ell \propto Y_d^T \propto \begin{pmatrix} \varepsilon^4 & \varepsilon^3 & \varepsilon \\ \varepsilon^3 & \varepsilon^2 & 1 \\ \varepsilon^3 & \varepsilon^2 & 1 \end{pmatrix}, \quad Y_\nu \propto \begin{pmatrix} \varepsilon^{|1\pm n_1|} & \varepsilon^{|1\pm n_2|} & \varepsilon^{|1\pm n_3|} \\ \varepsilon^{|n_1|} & \varepsilon^{|n_2|} & \varepsilon^{|n_3|} \\ \varepsilon^{|n_1|} & \varepsilon^{|n_2|} & \varepsilon^{|n_3|} \end{pmatrix}, \quad (4)$$

where $Y_{u,d,\ell,\nu}$ stand for the Yukawa couplings of quarks, charged leptons and neutrinos respectively, and n_i denote the U(1) charges of the heavy Majorana neutrinos. The heavy Majorana mass matrix is then given by

$$M_N \propto \begin{pmatrix} \varepsilon^{2|n_1|} & \varepsilon^{|n_1+n_2|} & \varepsilon^{|n_1+n_3|} \\ \varepsilon^{|n_1+n_2|} & \varepsilon^{2|n_2|} & \varepsilon^{|n_2+n_3|} \\ \varepsilon^{|n_1+n_3|} & \varepsilon^{|n_2+n_3|} & \varepsilon^{2|n_3|} \end{pmatrix}. \quad (5)$$

There is no unique choice for the right-handed neutrino charges n_1, n_2, n_3 , and several choices may be compatible with the low-energy neutrino data. We know, however, that the neutrino masses and mixing angles are related to the $\nu_L \nu_L$ contributions in the effective neutrino mass matrix

$$m_{eff} \approx m_\nu^D \frac{1}{M_N} m_\nu^{DT}, \quad (6)$$

which, if calculated from the charges in (4), is of the form:

$$m_{eff} \propto \begin{pmatrix} \varepsilon^2 & \varepsilon & \varepsilon \\ \varepsilon & 1 & 1 \\ \varepsilon & 1 & 1 \end{pmatrix}. \quad (7)$$

This form of m_{eff} is quite natural in the simplest see-saw models with a single expansion parameter and generic structures for the heavy and light Majorana mass matrices, due to cancellations that eliminate the dependences on the right-handed charges. Its predictions have been extensively analysed from a phenomenological point of view [44], and give a reasonable match to the data, provided there are no cancellations with large mixing in the charged lepton sector. Among other predictions, θ_{13} turns out to be naturally of the correct order of magnitude. It is interesting to note also that, to lowest order in ε , $Y_\ell Y_\ell^\dagger$ has the same structure as m_{eff} , namely

$$Y_\ell Y_\ell^\dagger \propto m_{eff} \propto \begin{pmatrix} \varepsilon^2 & \varepsilon & \varepsilon \\ \varepsilon & 1 & 1 \\ \varepsilon & 1 & 1 \end{pmatrix}. \quad (8)$$

The flavour mixing matrices are determined by the following diagonalizations of the Dirac and Majorana mass matrices:

$$V_\ell^T (Y_\ell Y_\ell^\dagger) V_\ell^* = \text{diag}(y_e^2, y_\mu^2, y_\tau^2), \quad (9)$$

$$V_D^T (Y_\nu Y_\nu^\dagger) V_D^* = \text{diag}(y_{\nu_1}^2, y_{\nu_2}^2, y_{\nu_3}^2), \quad (10)$$

$$U_N^T M_N U_N = \text{diag}(M_1, M_2, M_3), \quad (11)$$

$$U_\nu^T m_{eff} U_\nu = \text{diag}(m_{\nu_1}, m_{\nu_2}, m_{\nu_3}). \quad (12)$$

In terms of the above matrices, the Maki-Nakagawa-Sakata (MNS) matrix is given by

$$U_{MNS} \equiv U = V_\ell^\dagger U_\nu, \quad (13)$$

and can be parametrized as:

$$U = V \cdot \text{diag}(e^{-i\phi_1/2}, e^{-i\phi_2/2}, 1), \quad (14)$$

where

$$V = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - s_{23}s_{13}c_{12}e^{i\delta} & c_{23}c_{12} - s_{23}s_{13}s_{12}e^{i\delta} & s_{23}c_{13} \\ s_{23}s_{12} - c_{23}s_{13}c_{12}e^{i\delta} & -s_{23}c_{12} - c_{23}s_{13}s_{12}e^{i\delta} & c_{23}c_{13} \end{pmatrix}, \quad (15)$$

and c_{ij} and s_{ij} stand for $\cos \theta_{ij}$ and $\sin \theta_{ij}$, respectively.

2.1 Predictions for Neutrino Observables

Within the above framework of Abelian flavour symmetries there are ambiguities in the choices of coefficients, limited to a certain extent by requiring consistency with the experimental data. Here we proceed as follows: we take the expansion parameter to be $\varepsilon = 0.2$ and multiply the entries of Y_ℓ , Y_ν and M_N given by Eqs. (4) and (5) by coefficients in the range $\pm[0.6, 2]$ that we indicate by ℓ_{ij} , ν_{ij} and N_{ij} for Y_ℓ , Y_ν and M_N , respectively. Given our ignorance of the CP-violating phase δ , we consider the case $\delta = 0$ in the following: our results do not depend strongly on this choice.

We analyse $\mathcal{O}(10^9)$ combinations of these coefficients, as follows:

- (i) We select the charged lepton Yukawa coupling matrix Y_ℓ so that the correct charged-lepton mass hierarchies are reproduced, namely

$$\frac{m_\mu}{m_\tau} \sim 0.06 \quad \frac{m_e}{m_\tau} \sim 2.5 \times 10^{-3}. \quad (16)$$

- (ii) The coefficients ν_{ij} and N_{ij} of Y_ν and M_N are then matched to a light neutrino mass matrix m_{eff} of the form (7), with entries that deviate by a factor between 1/2 and 2 from those in Eq. (7).




Figure 1: *Correlations between the neutrino mixing angles before and after imposing the constraints on the model coefficients as discussed in the text. The solid lines indicate the experimental bounds, and the small black crosses represent models satisfying all the constraints. The two large magenta crosses correspond to benchmark models that are discussed in the text and used for numerical calculations.*

- (iii) We assume a neutrino mass hierarchy with a mass splitting of the order of the neutrino masses and $m_{\nu_3} \sim 0.05$ eV.
- (iv) We require the following ranges for the mixing angles [7]:

$$\begin{aligned}
 0.27 &< \sin^2 \theta_{12} < 0.35, \\
 0.34 &< \sin^2 \theta_{23} < 0.67, \\
 0.018 &< \sin^2 \theta_{13} < 0.033.
 \end{aligned} \tag{17}$$

The range on θ_{13} is consistent with the values reported by both [10, 11] at the $3 - \sigma$ level ¹.

- (v) We make a further selection by requiring that the hierarchy of eigenvalues of $Y_\ell Y_\ell^\dagger$ (which, as discussed above, has a similar structure to m_{eff} and $Y_\nu Y_\nu^\dagger$) preserves the order of the gauge eigenstates. This simply reduces the density of solutions in the plots, since the predictions for the neutrino parameters are similar also for the other cases.

In Fig. 1 we present the predictions for the neutrino mixing angles corresponding to the above criteria. We can see that, within this class of models, most solutions that reproduce the correct range of θ_{12} and θ_{23} also predict a neutrino mixing angle θ_{13} that is compatible with the data

¹Different choices of expansion parameters and coefficients lead to similar phenomenology. In ref. [39], an expansion parameter $\eta = 0.06$ and coefficients in the range $[\frac{1}{\sqrt{10}}, \sqrt{10}]$ are used.

Fit	Y_ℓ	Y_ν	M_N
1	$\begin{pmatrix} \varepsilon^4 & 1.5\varepsilon^3 & 2\varepsilon \\ 0.5\varepsilon^3 & -1.9\varepsilon^2 & -0.5\varepsilon \\ 2\varepsilon^3 & -1.9\varepsilon^2 & 1.5 \end{pmatrix}$	$\begin{pmatrix} \varepsilon^{ 1\pm n_1 } & \varepsilon^{ 1\pm n_2 } & 1.75\varepsilon^{ 1\pm n_3 } \\ \varepsilon^{ n_1 } & \varepsilon^{ n_2 } & 1.25\varepsilon^{ n_3 } \\ 1.3\varepsilon^{ n_1 } & \varepsilon^{ n_2 } & -2\varepsilon^{ n_3 } \end{pmatrix}$	$\begin{pmatrix} \varepsilon^{2 n_1 } & \varepsilon^{ n_1+n_2 } & -1.3\varepsilon^{ n_1+n_3 } \\ \varepsilon^{ n_1+n_2 } & \varepsilon^{2 n_2 } & \varepsilon^{ n_2+n_3 } \\ -1.3\varepsilon^{ n_1+n_3 } & \varepsilon^{ n_2+n_3 } & -\varepsilon^{2 n_3 } \end{pmatrix}$
	$\sin^2 \theta_{13}=0.027$	$\sin^2 \theta_{12}=0.30$	$\sin^2 \theta_{23}=0.64$
2	$\begin{pmatrix} \varepsilon^4 & 2\varepsilon^3 & -1.75\varepsilon \\ -0.5\varepsilon^3 & 1.9\varepsilon^2 & 0.5 \\ -0.5\varepsilon^3 & -0.7\varepsilon^2 & 1.25 \end{pmatrix}$	$\begin{pmatrix} \varepsilon^{ 1\pm n_1 } & \varepsilon^{ 1\pm n_2 } & 2\varepsilon^{ 1\pm n_3 } \\ 0.75\varepsilon^{ n_1 } & \varepsilon^{ n_2 } & -0.5\varepsilon^{ n_3 } \\ \varepsilon^{ n_1 } & \varepsilon^{ n_2 } & 1.25\varepsilon^{ n_3 } \end{pmatrix}$	$\begin{pmatrix} \varepsilon^{2 n_1 } & \varepsilon^{ n_1+n_2 } & -\varepsilon^{ n_1+n_3 } \\ \varepsilon^{ n_1+n_2 } & \varepsilon^{2 n_2 } & \varepsilon^{ n_2+n_3 } \\ -\varepsilon^{ n_1+n_3 } & \varepsilon^{ n_2+n_3 } & -\varepsilon^{2 n_3 } \end{pmatrix}$
	$\sin^2 \theta_{13}=0.019$	$\sin^2 \theta_{12}=0.28$	$\sin^2 \theta_{23}=0.40$

Table 1: *Indicative textures for Y_ℓ , Y_ν and M_N , corresponding to the two crosses in Fig 1, with the indicated values of the neutrino mixing angles. The n_i are Abelian charges to be determined.*

from [10, 11]. Furthermore, one can see that, within the narrow range of θ_{13} indicated by the measurements of [10, 11], the model predictions are in the mid-lower range of θ_{12} . On the other hand, in the case of θ_{23} , solutions are found in two disconnected regions in the plane $(\sin^2 \theta_{13}, \sin^2 \theta_{23})$, at either end of the range that is in agreement with a global statistical analysis of the data performed in [45]. However, we did not find any correlation between the preferred values of θ_{23} and θ_{12} in Fig. 1, since the points in either of the two favoured regions of the right panel are distributed homogeneously in the left panel.

We would like to point out that the fits in Fig. 1 are independent of the charges n_i , which affect Y_ν and M_N but not their combination in m_{eff} . On the other hand, the choices of the n_i do affect the rates for charged LFV processes, whose magnitudes provide important inputs on the right-handed sector of the neutrino mass matrices. Charged LFV processes are therefore powerful probes of parameters that cannot be constrained by lepton hierarchies or by better measurements of the neutrino parameters. We return to this issue later in the paper.

As benchmarks for studying LFV in subsequent sections, we have identified two representative sets of textures from the many points that fit the neutrino data. These are indicated with crosses in Fig. 1. Since we are working with an Abelian flavour symmetry, we focus on solutions where lepton mixing arises from large off-diagonal elements in V_ℓ (in the basis of the GUT multiplets). In this case, the observed mixing is directly generated by the flavour symmetry - since it is correlated with the flavour charges of the charged leptons - and does not arise purely from the see-saw conditions, which often require more fine tuning.

These benchmarks are chosen as representatives of the two different regions identified in Fig 1: Fit 1 lies in the right region of the right panel of Fig 1, with larger θ_{23} , and Fit 2 lies in the left region of this panel, with smaller θ_{23} . Their parameters are listed in Table 1. For the reason discussed above, the mixing angles are insensitive to the values of the charges n_i , which determine the levels of the LFV processes, as we see in the following sections.

3 Charged-Lepton-Flavour Violation

We work in the constrained minimal supersymmetric standard model (CMSSM) with universal scalar, gaugino masses and trilinear terms at the GUT scale, m_0 , $M_{1/2}$ and A_0 , respectively. The physical values of these masses are obtained by integrating the renormalization-group equations (RGEs) of the MSSM supplemented with right-handed neutrinos. Since the Dirac neutrino and charged-lepton Yukawa couplings cannot, in general, be diagonalized simultaneously, and since both lepton Yukawa couplings appear in the RGEs, the lepton Yukawa matrices and the slepton mass matrices at low energies cannot be diagonalized simultaneously, either.

In the basis where the charged-lepton masses m_ℓ are diagonal, the soft slepton-mass matrix acquires a correction that contains off-diagonal contributions from renormalization at scales below M_{GUT} , of the form [32]:

$$\delta m_\ell^2 \propto \frac{1}{16\pi^2} (6m_0^2 + 2A_0^2) Y_\nu^\dagger Y_\nu \ln\left(\frac{M_{GUT}}{M_N}\right), \quad (18)$$

where M_N is the intermediate scale where the effective neutrino-mass operator is formed. The physical charged slepton masses are obtained by numerical diagonalization of the following 6×6 matrix:

$$m_\ell^2 = \begin{pmatrix} m_{LL}^2 & m_{LR}^2 \\ m_{RL}^2 & m_{RR}^2 \end{pmatrix}, \quad (19)$$

where all the entries are 3×3 matrices in flavour space. Using the basis where Y_ℓ is diagonal, it is convenient to write the 3×3 entries of (19) in the forms:

$$m_{LL}^2 = (m_{\tilde{\ell}}^{\text{diag}})^2 + \delta m_\ell^2 + m_\ell^2 - \frac{1}{2}(2M_W^2 - M_Z^2) \cos 2\beta, \quad (20)$$

$$m_{RR}^2 = (m_{\tilde{\ell}_R}^{\text{diag}})^2 + m_\ell^2 - (M_Z^2 - M_W^2) \cos 2\beta, \quad (21)$$

$$m_{RL}^2 = (A_\ell^{\text{diag}} + \delta A_\ell - \mu \tan \beta) m_\ell, \quad (22)$$

$$m_{LR}^2 = m_{RL}^{2\dagger}. \quad (23)$$

Here $\tan \beta$ is the standard ratio of the two MSSM Higgs vevs, $(m_{\tilde{\ell}}^{\text{diag}})^2$, $(m_{\tilde{\ell}_R}^{\text{diag}})^2$ and A_ℓ^{diag} denote the diagonal contributions to the corresponding matrices, obtained by numerical integration of the renormalization-group equations, and δm_ℓ^2 and δA_ℓ denote the corrections with off-diagonal terms that appear because Y_ν and Y_ℓ cannot be diagonalized simultaneously.

The full mass matrix for left- and right-handed sneutrinos has a 12×12 structure, given in terms of 3×3 Dirac, Majorana and sneutrino mass matrices. The effective 3×3 mass-squared matrix for the left-handed sneutrinos has the same form as the m_{LL}^2 part (23) of the 6×6 charged-slepton matrix (19), with the difference that now the Dirac masses are absent. In Ref. [33] it was shown that is sufficient to use

$$m_\nu^2 = (m_{\tilde{\ell}}^{\text{diag}})^2 + \delta m_\ell^2 + \frac{1}{2} M_Z^2 \cos 2\beta. \quad (24)$$

The matrix responsible for LFV is $V_{LFV} = V_D^\dagger V_\ell$, and the slepton mass matrices contain off-diagonal terms generated by:

$$m_{LL}^2 = V_{LFV}^\dagger (m_{LL}^2)_{\text{diag}} V_{LFV}, \quad (25)$$

while the A -terms become:

$$A_\ell = V_{LFV}^T (A_\ell)_{\text{diag}}. \quad (26)$$

Here $(m_{LL}^2)_{\text{diag}}$ and $(A_\ell)_{\text{diag}}$ are the soft terms resulting from the RGE running of the universal soft terms at the GUT scale.

The matrix element of the electromagnetic-current operator between two distinct lepton mass eigenstates ℓ_i and ℓ_j is given in general by

$$\begin{aligned} \mathcal{T}_\lambda &= \langle \ell_i | (p - q) | \mathcal{J}_\lambda | \ell_j(p) \rangle \\ &= \bar{u}_i(p - q) [m_j i \sigma_{\lambda\beta} q^\beta (A_M^L P_L + A_M^R P_R) + \\ &\quad (q^2 \gamma_\lambda - q_\lambda \gamma \cdot q) (A_E^L P_L + A_E^R P_R)] u_j(p), \end{aligned} \quad (27)$$

where q is the photon momentum. The coefficients A_M and A_E denote contributions from neutralino/charged-slepton and chargino/sneutrino exchanges. The amplitude of the LFV process is then proportional to $\mathcal{T}_\lambda \epsilon^\lambda$, where ϵ^λ is the photon-polarization vector. The branching ratios (BRs) of the decays $\ell_j \rightarrow \ell_i + \gamma$ are calculated using the formulae of Ref. [32].

3.1 Predictions for Radiative Decays and LFV at LHC

We now study the conditions under which the favoured ranges of lepton masses and of neutrino mixing, can lead in a natural way to observable signatures for charged LFV. As already discussed, while the neutrino parameters are independent of the charges n_i , this is not the case for LFV. As a result, LFV can provide a way to probe the right-handed neutrino sector, for which only limited information is available.

In a previous section, we identified two representative benchmark fits suitable for studying the expected rates for charged LFV. As we saw previously, the level of charged LFV is determined by the product $V_{LFV} = V_D^\dagger \cdot V_\ell$, and thus by the charges n_i which enter in the Dirac neutrino mixing matrix $V_D^\dagger$. However, in general, different choices of n_i lead to different solutions for a large V_ℓ in the GUT eigenstate basis, and it is possible to group these solutions in the following categories:

- Solutions with a large V_ℓ lead to large LFV, for all charge combinations that generate a small $V_D^\dagger$.
- For solutions with large $V_D^\dagger$, the mixing may add to or subtract from that of the leptons, depending on coefficients and phases. In the first case, large LFV is expected, whereas in the second case LFV is suppressed. However, it is difficult to find suppression in all three relevant mixing angles simultaneously.

n_i	(i) $\{n_1 = 1, n_2 = 0, n_3 = 0\}$	(ii) $\{n_1 = 2, n_2 = 1, n_3 = 0\}$	(iii) $\{n_1 = 2, n_2 = 0, n_3 = 1\}$	(iv) $\{n_1 = 0, n_2 = 1, n_3 = 0\}$
V_{LFV}	$\begin{pmatrix} 0.805 & -0.385 & -0.451 \\ 0.182 & 0.885 & -0.429 \\ 0.565 & 0.263 & 0.782 \end{pmatrix}$	$\begin{pmatrix} 0.805 & -0.385 & -0.452 \\ -0.064 & 0.700 & -0.711 \\ 0.590 & 0.601 & 0.539 \end{pmatrix}$	$\begin{pmatrix} -0.820 & 0.343 & 0.458 \\ 0.494 & 0.829 & 0.263 \\ -0.289 & 0.441 & -0.849 \end{pmatrix}$	$\begin{pmatrix} 0.806 & -0.401 & -0.436 \\ -0.437 & -0.899 & 0.016 \\ -0.399 & 0.178 & -0.901 \end{pmatrix}$

Table 2: Values of V_{LFV} with $\varepsilon = 0.2$ for Fit 2 of Table 1 with large V_ℓ . From left to right, we see the following features: (i) V_D is small, leading to large V_{LFV} , (ii) and (iii) V_D is large and so is V_{LFV} , (iv) V_D is large and cancellations with V_ℓ occur in the 2-3 sector, suppressing LFV. In the last two examples, the heavy Majorana mass matrix has an inverted hierarchy.

These observations indicate that, even before looking at the influence of the charges n_i or performing specific computations, LFV processes can be quite large for natural choices of textures.

A survey of the dependences of the entries in V_{LFV} on the different right-handed neutrino charges is given in Table 2. For simplicity, we focus on Fit 2, noting that similar results hold for Fit 1. We see explicitly that different values of the charges n_i result in different levels of LFV, even though the m_{eff} eigenvalues are unchanged, thanks to the see-saw conditions.

In addition to the flavour-mixing effects, the rates for flavour non-conserving decays are sensitive to other physical quantities. In the supersymmetric GUT context explored here, they depend on the masses of the sparticles that mediate the flavour non-conserving processes. In the CMSSM studied here, we parametrize these masses in terms of the universal GUT-scale parameters m_0 and $M_{1/2}$, and use the RGEs of the MSSM to calculate the low-energy sparticle masses, taking into account low-energy threshold effects. Other relevant MSSM free parameters are the trilinear coupling A , the sign of the Higgs mixing parameter μ , and the value of $\tan\beta$. Here we choose values of the CMSSM input parameters that are consistent with cosmological relic density constraints, as well as LHC results.

The recent LHC measurement of the Higgs mass [46, 47] imposes severe constraints in the CMSSM parameter space. Specifically, a Higgs masses $m_h \sim 125$ GeV implies, in general, a relatively heavy sparticle spectrum that is consistent with the cosmological constraint on the neutralino relic density only in limited regions. A global analysis of the CMSSM parameter space was performed in [41], and yielded two almost equally good fits to the available data, one with relatively low sparticle masses and $\tan\beta \sim 16$, and the other with larger sparticle masses and $\tan\beta \sim 45$, corresponding to the following CMSSM sets of parameters:

$$\begin{aligned}
(a) \quad & \tan\beta = 16, \quad m_0 = 300 \text{ GeV}, \quad M_{1/2} = 910 \text{ GeV}, \quad A_0 = 1320 \text{ GeV}, \\
(b) \quad & \tan\beta = 45, \quad m_0 = 1070 \text{ GeV}, \quad M_{1/2} = 1890 \text{ GeV}, \quad A_0 = 1020 \text{ GeV}.
\end{aligned} \tag{28}$$

Both these choice of input parameters are compatible with m_h measurements, within theoretical uncertainties [41].

Point (a) predicts a lighter sparticle spectrum, which generally favours larger LFV rates at




Figure 2: Predictions for the rare LFV decays $\ell_i \rightarrow \ell_j \gamma$ as a function of the heaviest right-handed neutrino mass M_N , for the benchmark point displayed in (28) (a) (thick line), (b) (thin line) and using some of the neutrino mixing fits shown in Table 2. The solid lines correspond to cases (iv) and dashed lines (iii). The horizontal solid lines indicate the current experimental upper bounds.

accelerators, since there is more phase space at the LHC and smaller energies are required for sparticle production at a LC. On the other hand, point (b) is indicative of models with a heavier spectrum, presenting greater experimental challenges. Regarding cosmological considerations, point (a) belongs to the region where the WMAP-favoured range of $\Omega_\chi h^2$ is achieved via $\chi - \tilde{\tau}$ coannihilation², whereas point (b) lies in the funnel region where the neutralino LSP annihilates rapidly via direct-channel H/A poles.

²We note that in this region the $\chi - \tilde{\tau}$ mass difference is very small, offering other experimental challenges and opportunities [48, 49].

In Fig. 2 we show numerical predictions for the LFV branching ratios arising from the textures introduced above (which match the neutrino data) for the CMSSM parameters specified in (28) (which are compatible with the LHC constraints). We show the effect of varying M_N from 6×10^{14} GeV down to 10^{12} GeV, for Fit 2 of Table 1 and for two choices of right-handed neutrino charges, n_i (0,1,0) and (2,0,1) respectively). We can see that the experimental upper bounds on $\text{BR}(\mu \rightarrow e\gamma)$ can be reached for suitable values of the Yukawa structure, even with the heavy sparticle spectrum suggested by the higher-mass benchmark (b).

It is clear from the three panels of Fig. 2 that the combined uncertainties from the unknown right-handed charges n_i and the unknown model coefficients inevitably introduce significant uncertainties. On the other hand, the rates for radiative charged LFV exhibit different correlations in the different benchmark scenarios, suggesting that future observations could possibly provide important input into the selection of model parameters. Further information could be obtained by LFV studies at a Linear Collider (LC), as discussed in the next Section. In general, we find that:

- The branching ratios are larger for the lower-mass scenario with $\tan\beta = 16$, due to the lighter spectrum.
- Large values of A_0 may induce some cancellations in the branching ratios for radiative decays, suppressing them strongly. In such regions, colliders provide the best chances for probing LFV. We discuss some examples at the end of the next Sections.
- A lighter spectrum implies smaller threshold energies and larger cross sections for slepton production at a LC.

In previous work [26], we concluded that the bounds of $\text{BR}(\tau \rightarrow \mu\gamma)$ disfavour mixings that could lead to observable signals at the LHC in the LL sector, whereas significantly larger mixing were still possible in the RR sector. This occurs for a broad region of points like scenario (b), where the size of A_0 is limited by electroweak symmetry breaking and $m_h \sim 125$ GeV is realized with a heavy spectrum with large $\tan\beta$. Point (a), on the other hand, has a light spectrum with large trilinear terms that are nevertheless compatible with electroweak symmetry breaking. As a result, LR flavour mixing in the charged slepton masses may induce non-trivial LFV in slepton decays.

In what follows, we revisit the $\chi_2 \rightarrow \chi + \tau^\pm \mu^\mp$ decay mode for the benchmark points (a) and (b), in order to determine whether the relevant ratios

$$R_{\tau\mu} = \Gamma(\chi_2 \rightarrow \chi + \tau^\pm + \mu^\mp) / \Gamma(\chi_2 \rightarrow \chi + \tau^\pm + \tau^\mp) \quad (29)$$

can be of the order of 10%, as required for a signal that can be distinguished from the background [24, 26]. We include all the contributing on-shell sfermion exchange diagrams, as given in [50]:

$$\text{BR}(\chi_2 \rightarrow \chi \tau^\pm \mu^\mp) = \sum_{i=1}^3 \left[\text{BR}(\chi_2 \rightarrow \tilde{\ell}_i \mu) \text{BR}(\tilde{\ell}_i \rightarrow \tau \chi) + \text{BR}(\chi_2 \rightarrow \tilde{\ell}_i \tau) \text{BR}(\tilde{\ell}_i \rightarrow \mu \chi) \right]. \quad (30)$$




Figure 3: The ratio defined in Eq. (29) for the CMSSM points (a) (thick line) and (b) (thin line), Eq. 28, using the mixing options (iv) (solid lines) and (iii) (dashed lines) of Table 2.

We present in the left panel of Fig. 3 the ratio of the flavor-violating decay width $\Gamma(\chi_2 \rightarrow \chi + \tau^\pm + \mu^\mp)$ to the CMSSM flavor-conserving decay width $\Gamma(\chi_2 \rightarrow \chi + \tau^\pm + \tau^\mp)$. We see that for point (a) predictions are within the reach of the LHC, whereas for point (b) the predictions are below the expected experimental sensitivity for LFV observation in neutralino decays at the LHC.

4 LFV Observation at a Linear Collider

In previous work [21], we studied the different LFV channels at a LC, focusing on the production of $\mu^\pm\tau^\pm$ pairs. In supersymmetric models where LFV is produced by lepton-slepton vertices, observable signatures may occur either directly, in slepton-pair production, or indirectly, via slepton production in cascade decays [16]. If the flavour mixing is introduced in the LL slepton sector, as is the case for the models under consideration here, pair production dominates in slepton-pair production and LFV decays such as:

$$\begin{aligned} e^+e^- &\rightarrow \tilde{\ell}_i^-\tilde{\ell}_j^+ \rightarrow \tau^\pm\mu^\mp\tilde{\chi}_1^0\tilde{\chi}_1^0, \\ e^+e^- &\rightarrow \tilde{\nu}_i\tilde{\nu}_j^c \rightarrow \tau^\pm\mu^\mp\tilde{\chi}_1^+\tilde{\chi}_1^-. \end{aligned} \quad (31)$$

At the CMSSM benchmark point introduced above, which favoured by current experimental data and cosmological observations, the channel mediated by charged sleptons clearly dominates over the sneutrino-pair production process, and may lead to a cross section of the order of 1 fb, the reference value used in [21], for a LFV signal of $\mu^\pm\tau^\pm$ pairs that can be distinguished from the background, according to the considerations in [19]. Here, we extend our previous results by considering the full structure of the Yukawa couplings, thus comparing the LFV production

of charged leptons of all generations. In addition, we update the supersymmetric benchmark points used previously, choosing a new CMSSM benchmark scenario compatible with the LHC measurement of m_h .




Figure 4: Values of the cross sections $\sigma(e^+e^- \rightarrow \tilde{\ell}_i^-\tilde{\ell}_j^+ \rightarrow \ell_a^\pm\ell_b^\mp + 2\chi^0)$ ($\ell_a \neq \ell_b$ as indicated for each panel) as functions of $\sqrt{s}$ for $M_N = 10^{14}$ GeV. The line styles as the same as those in Fig. 2.

We present in Fig. 4 the expected cross sections as functions of $\sqrt{s}$ for Fits 1 and 2 of Table 1 and the same sets of charges as those in Fig. 2. For definiteness, the right-handed neutrino mass scale is set to $M_N = 10^{14}$ GeV. We observe that, by varying this scale in a similar range as in Fig. 2, the cross sections also change by approximately an order of magnitude: according to the see-saw conditions, larger values of M_N imply larger Dirac neutrino Yukawa couplings, and thus stronger RGE running. The results depend sensitively on the particle spectrum. If




Figure 5: *Upper panels: branching ratios for the CMSSM points (a) (thick line) and (b) (thin line), Eq. (28), using case (ii) the neutrino mixing fits shown in Table 2. The lower panels show the corresponding prediction for the LFV cross sections.*

we restrict ourselves to the benchmark CMSSM parameter choice discussed above, we would require $\sqrt{s}$ beyond 3 TeV in order to have significant LFV signals in slepton mixing. However, once above threshold, the uncertainties in the choices of right-handed neutrino charges and unknown model coefficients have similar impacts on the LC LFV cross sections to those on the radiative decays discussed in the previous section.

In the case of point (a) we find cancellations in the branching ratios for some of the neutrino mixing fits. This happens because of the large value of A_0 , which leads to significant cancellations among the different amplitudes that contribute to the LFV decay amplitudes [32, 33, 34]. Using fits (ii) and (iv) from Table 2 and the benchmark point (a), we find this type of cancel-




Figure 6: *The ratio defined in Eq. (29) for the CMSSM points (a) (thick line) and (b) (thin line), Eq. (28), using the mixing options (ii) of Table 2.*

lation in $\text{BR}(\tau \rightarrow \mu\gamma)$ and $\text{BR}(\tau \rightarrow e\gamma)$ for a range of values of the right-handed neutrino scale M_N , as shown in the upper panels of Fig. 5. The lower panels show the corresponding predictions for LFV stau decays at a LC when $M_N = 2 \times 10^{13}$ GeV. We see that the cancellations in the branching ratios do not occur for the LFV LC signals. This implies that it is possible to observe slepton flavor oscillations at the LC, in cases where LFV would be undetectable in rare charged lepton decays.

Fits (i) and (ii) in Table 2 correspond to cases with larger mixing in the matrix V_{LFV} . In these two cases, the choice of charges induces an inverse hierarchy in M_N . In both cases, the large A -terms and large 2-3 mixing make cancellations possible. We display results for case (i) but not for case (ii), which is quite similar.

5 Conclusions

Abelian flavour symmetries provide interesting possibilities for understanding the hierarchy of fermion masses and mixing. Despite uncertainties in the choice of $\mathcal{O}(1)$ coefficients, they offer useful insight into physical observables, and provide specific predictions for the signals to be expected in various detection channels, which serve as diagnostic tools for discriminating between different models.

In our work, we have explored these issues using updated experimental input from neutrino data, particularly recent measurements of θ_{13} , MEG and the LHC. We have revisited the signatures of charged LFV in theoretically-motivated scenarios, studying the correlations arising in CMSSM models with parameter values that are favoured by the LHC and cosmological

considerations. Because of their sensitivity to flavour symmetries and model parameters that are not constrained by the neutrino data, particularly those linked to the right-handed neutrino sector, LFV searches may become a powerful tool for distinguishing between different theoretical scenarios.

Within an $SU(5)$ GUT framework supplemented by an Abelian flavour symmetry, we first performed a scan of different fits to the neutrino data, paying attention to the naturalness of the fit and avoiding artificial cancellations arising from specific choices of coefficients. This leads to a pattern of correlations between the neutrino mixing angles that are compatible with the global analysis of neutrino data in [45].

We then looked at the expectations for LFV processes in the above models, identifying the ranges of parameters where observable signatures are possible. In general, fits with similar predictions for the neutrino parameters may lead to different LFV predictions. However, the recent input on θ_{13} , combined with the new MEG bound on $\mu \rightarrow e\gamma$ as well as LHC data, do constrain the allowed structures. Further precision in the determination of neutrino parameters could lead to restrictions on the choices of model coefficients, but would not constrain the right-handed neutrino charges. However, new input in this respect could be provided by the rates for LFV processes, since their magnitudes are directly linked to these charges, unlike the neutrino mass and mixing parameters.

In the cases we studied, it was possible to establish correlations between the expected rates for radiative LFV decays, LFV processes at the LHC and $\tau\mu$ pair production at a future LC, for different possibilities for the structure of the heavy Majorana neutrino masses. As we have shown, extending previous studies by including the full 3×3 neutrino mixing leads to interesting correlations between the LFV production of charged leptons of different generations at a LC. However, within the CMSSM framework studied, the absence of a supersymmetry signal in the LHC data and the discovery of a neutral Higgs weighing ~ 125 GeV imply that observation of slepton flavour violation at a LC will be possible for only energies beyond 1 TeV, in models with large Dirac neutrino couplings and a heavy Majorana neutrino scale M_N . On the other hand, scenarios less constrained than the CMSSM that could fit m_h with a lighter sparticle spectrum might predict observable LFV signals at smaller energies.

Acknowledgements The work of J.E. is supported partly by the London Centre for Terauniverse Studies (LCTS), using funding from the European Research Council via the Advanced Investigator Grant 267352. S.L. thanks the CERN Theory Division for its kind hospitality. The work of M. C. is supported by MultiDark under Grant No. CSD2009-00064 of the Spanish MICINN Consolider-Ingenio 2010 Program. M. E. G. and M. C. acknowledge further support from the MICINN project FPA2011-23781, from the Grant MICINN-INFN(PG21)AIC-D-2011-0724 and the project P07FQM02962 funded by “Junta de Andalucía”.

References

- [1] Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **81** (1998) 1562.
- [2] Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **82** (1999) 1810; Phys. Rev. Lett. **82** (1999) 2430; Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **87** (2001) 071301.
- [3] K. Eguchi *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **90** (2003) 021802; T. Araki *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **94** (2005) 081801.
- [4] M.H. Ahn *et al.* [K2K Collaboration], Phys. Rev. Lett. **90** (2003) 041801.
- [5] D.G. Michael *et al.* [MINOS Collaboration], Phys. Rev. Lett. **97** (2006) 191801.
- [6] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, Phys. Rev. D **84** (2011) 053007.
- [7] M. C. González-García, M. Maltoni, J. Salvado and T. Schwetz, arXiv:1209.3023 [hep-ph].
- [8] K. Abe *et al.* [T2K Collaboration], Phys. Rev. Lett. **107** (2011) 041801.
- [9] P. Adamson *et al.* [MINOS Collaboration], Phys. Rev. Lett. **107** (2011) 181802.
- [10] F. P. An *et al.* [Daya Bay Collaboration], Phys. Rev. Lett. **108** (2012) 171803.
- [11] J. K. Ahn *et al.* [RENO Collaboration], Phys. Rev. Lett. **108** (2012) 191802.
- [12] Y. Abe *et al.* [Double Chooz Collaboration], arXiv:1301.2948 [hep-ex].
- [13] For reviews, see:
M. Raidal *et al.*, Eur. Phys. J. C **57** (2008) 13; Y. Kuno and Y. Okada, Rev. Mod. Phys. **73** (2001) 151.
- [14] J. Adam *et al.* [MEG Collaboration], Phys. Rev. Lett. **107** (2011) 171801.
- [15] K. Nakamura *et al.* [Particle Data Group], J. Phys. G **37** (2010) 075021.
- [16] N. Arkani-Hamed, H. Cheng, J. L. Feng and L. J. Hall, Phys. Rev. Lett. **77** (1996) 1937; Nucl. Phys. B **505** (1997) 3.
- [17] J. Hisano, M. M. Nojiri, Y. Shimizu and M. Tanaka, Phys. Rev. D **60**, 055008 (1999); M. Guchait, J. Kalinowski and P. Roy, Eur. Phys. J. C **21**, 163 (2001).
- [18] F. Deppisch, J. Kalinowski, H. Päs, A. Redelbach and R. Rückl, arXiv:hep-ph/0401243, work performed for the LCH study group.
- [19] F. Deppisch, H. Päs, A. Redelbach, R. Rückl and Y. Shimizu, Phys. Rev. D **69** (2004) 054014.

- [20] M. Cannoni, S. Kolb and O. Panella, Phys. Rev. D **68** (2003) 096002 [hep-ph/0306170]; M. Cannoni, C. Carimalo, W. Da Silva and O. Panella, Phys. Rev. D **72** (2005) 115004 [Erratum-ibid. D **72** (2005) 119907] [hep-ph/0508256]; M. Cannoni and O. Panella, Phys. Rev. D **79** (2009) 056001 [arXiv:0812.2875 [hep-ph]].
- [21] E. Carquin, J. Ellis, M. E. Gómez and S. Lola, JHEP **1111** (2011) 050.
- [22] A. Abada, A. J. R. Figueiredo, J. C. Romão and A. M. Teixeira, JHEP **1208** (2012) 138 [arXiv:1206.2306 [hep-ph]].
- [23] N. V. Krasnikov, JETP Lett. **65** (1997) 148; S. I. Bityukov and N. V. Krasnikov, arXiv:hep-ph/9806504, (*10th International Seminar on High-Energy Physics Suzdal, Russia*); K. Agashe and M. Graesser, Phys. Rev. D **61** (2000) 075008.
- [24] I. Hinchliffe and F. E. Paige, Phys. Rev. D **63** (2001) 115006;
- [25] J. Hisano, R. Kitano and M. M. Nojiri, Phys. Rev. D **65** (2002) 116002; D. Carvalho, J. Ellis, M. Gómez, S. Lola and J. Romão, Phys. Lett. B **618** (2005) 162.
- [26] E. Carquin, J. Ellis, M. E. Gómez, S. Lola and J. Rodríguez-Quintero, JHEP **0905** (2009) 026.
- [27] A. Abada, A. J. R. Figueiredo, J. C. Romão and A. M. Teixeira, JHEP **1108** (2011) 099 [arXiv:1104.3962 [hep-ph]].
- [28] J. N. Esteves, J. C. Romão, A. Villanova del Moral, M. Hirsch, J. W. F. Valle and W. Porod, JHEP **0905** (2009) 003 [arXiv:0903.1408 [hep-ph]].
- [29] M. Hirsch, W. Porod, F. Staub and C. Weiss, Phys. Rev. D **87**, 013010 (2013) [arXiv:1211.0289 [hep-ph]].
- [30] M. Gell-Mann, P. Ramond and R. Slansky, *Proceedings of the Stony Brook Supergravity Workshop, New York, 1979*, eds. P. Van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam).
- [31] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57** (1986) 961.
- [32] J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. D **53** (1996) 2442.
- [33] M. Gómez, G. Leontaris, S. Lola and J. Vergados, Phys. Rev. D **59** (1999) 116009.
- [34] J. R. Ellis, M. E. Gómez, G. K. Leontaris, S. Lola and D. V. Nanopoulos, Eur. Phys. J. C **14** (2000) 319.
- [35] S. Antusch, E. Arganda, M. J. Herrero and A. M. Teixeira, JHEP **0611** (2006) 090 [hep-ph/0607263].

- [36] J. Hisano, D. Nomura and T. Yanagida, Phys. Lett. B **437** (1998) 351; W. Buchmüller, D. Delepine and F. Vissani, Phys. Lett. B **459** (1999) 171; W. Buchmüller, D. Delepine and L. T. Handoko, Nucl. Phys. B **576** (2000) 445; J. L. Feng, Y. Nir and Y. Shadmi, Phys. Rev. D **61** (2000) 113005; J. Sato and K. Tobe, Phys. Rev. D **63** (2001) 116010; J. Hisano and K. Tobe, Phys. Lett. B **510** (2001) 197; S. Baek, T. Goto, Y. Okada and K. Okumura, Phys. Rev. D **64** (2001) 095001; S. Lavignac, I. Masina and C.A. Savoy, Phys. Lett. B **520** (2001) 269; D. Carvalho, J. Ellis, M. Gómez and S. Lola, Phys. Lett. B **515** (2001) 323; T. Blazek and S. F. King, Phys. Lett. B **518** (2001) 109.
- [37] S. Lola and G.G. Ross, Nucl. Phys. B **553** (1999) 81.
- [38] J. R. Ellis, M. E. Gómez and S. Lola, JHEP **0707** (2007) 052.
- [39] W. Buchmüller, V. Domcke and K. Schmitz, JHEP **1203** (2012) 008.
- [40] D. Meloni, JHEP **1110** (2011) 010
- [41] O. Buchmueller *et al.*, Eur. Phys. J. C **72** (2012) 2243 [arXiv:1207.7315 [hep-ph]].
- [42] See for instance some of the textures studied by G. Altarelli and F. Feruglio, Phys. Lett. B **439** (1998) 112 and JHEP **9811** (1998) 021.
- [43] M.E. Gómez, S. Lola, P. Naranjo and J. Rodríguez-Quintero, JHEP **0904** (2009) 043.
- [44] G. Altarelli and F. Feruglio, “Theoretical models of neutrino masses and mixings,” Springer Tracts Mod. Phys. **190** (2003) 169.
- [45] G.L. Fogli, E. Lisi, A. Marrone, D. Montanino and A. Palazzo, A.M. Rotunno, Phys. Rev. D **86** (2012) 013012.
- [46] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716** (2012) 1.
- [47] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716** (2012) 30.
- [48] M. Citron, J. Ellis, F. Luo, J. Marrouche, K. A. Olive and K. J. de Vries, arXiv:1212.2886 [hep-ph].
- [49] M. Cannoni, O. Panella, M. Pioppi and M. Santoni, Phys. Rev. D **86** (2012) 037702 [arXiv:1206.5759 [hep-ph]].
- [50] A. Bartl, K. Hidaka, K. Hohenwarter-Sodek, T. Kernreiter, W. Majerotto and W. Porod, Eur. Phys. J. C **46** (2006) 783 [arXiv:hep-ph/0510074].